



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

genera, while some of the species of these two groups are most certainly much more nearly related to each other than they are to any of the interposed groups. Our Robin group is allotted to *Turdus*, and forms the only American species of the genus, except *T. rufitorques* of Mexico and Central America.

It seems like returning to the 'good old times' to see such groups as the Mimidæ, Regulidæ, Paridæ, Certhiidæ, etc., installed again as full-fledged families.

*Parus* is restricted to a group of Old World Titmice, the American species hitherto referred to *Parus* being placed in *Pœcile* Kaup, except *P. gambeli*, for which the new genus *Pœcilodes* Bianchi (1902) is adopted.

The recent additions to the list of described forms are given at their face value, with, however, references to adverse opinions when any such have been made public. In short, the care, thoroughness and fairness of Dr. Sharpe's great work will long render it a most invaluable aid to every systematic ornithologist.—J. A. A.

**Ridgway on New American Birds.**—Mr. Ridgway, in preparing Part III of his 'Birds of North and Middle America,' has found it desirable to describe a number of new genera, species, and subspecies.<sup>1</sup> The new genera comprise the following four genera of Swallows, as follows: *Alopochelidon*, type, *Hirundo fucata* Temm.; *Orochelidon*, type, *Petrochelidon murina* Cass.; *Diplochelidon*, type, *Hirundo melanoleuca* Wied; *Lamprochelidon*, type, *Hirundo euchrysea* Gosse. The new species and subspecies, 29 in number, are mostly from Mexico and Central America, but the following come within the scope of the A. O. U. Check-List: (1) *Budytes flavus alascensis*, Western Alaska; (2) *Vireo huttoni cognatus*, Cape district of Lower California; (3) *Vireo bellii arizonæ*, western Texas and Arizona; (4) *Lanius ludovicianus mearnsi*, San Clemente and Santa Margarita Islands, L. Cal.; (5) *Bæolophus inornatus restrictus*, vicinity of San Francisco Bay, Cal.; (6) *B. i. murinus*, northern Lower California; (7) *Psaltiriparus minimus saturatus*, Mount Vernon, Wash.; (8) *Chamæa fasciata rufula*, central coast region of California; (9) Mississippi Valley and Great Plains region, north to Alberta.—J. A. A.

**Nelson on New Birds from Mexico.**—The 13 new species and subspecies here described<sup>2</sup> were mainly collected by Messrs. Nelson and Goldman in southwestern Mexico during the winter of 1902-03. They

<sup>1</sup> Descriptions of New Genera, Species, and Subspecies of American Birds. Proc. Biol. Soc. Wash., XVI, pp. 105-113, Sept. 30, 1903.

Diagnoses of Nine New Forms of American Birds. *Ibid.*, pp. 167-170, Nov. 30, 1903.

<sup>2</sup> Descriptions of New Birds from Southern Mexico. By E. W. Nelson. Proc. Biol. Soc. Wash. XVI, pp. 151-160, Nov. 30, 1903.